

113. Aach RD, Stevens CE, Hollinger FB, et al. Hepatitis C virus infection in post-transfusion hepatitis: an analysis with first- and second-generation assays. *N Engl J Med* 1991;325:1325-9.
114. Sheridan RL. Evaluating and managing burn wounds. *Dermatol Nurs* 2000;12(1): 17-30.
115. Heimbach D, Engrav L, Grube B, Marvin J. Burn depth: a review. *World J Surg* 1992;16:10-5.
116. Anselmo VJ, Zawacki BE. Multispectral photographic analysis: a new quantitative tool to assist in the early diagnosis of thermal burn depth. *Ann Biomed Eng* 1977;5:179-93.
117. Afromowitz MA, Van Liew GS, Heimbach DM. Clinical evaluation of burn injuries using an optical reflectance technique. *IEEE Trans Biomed Eng* 1987;34:114-27.
118. Afromowitz MA, Callis JB, Heimbach DM, DeSoto LA, Norton MK. Multispectral imaging of burn wounds: a new clinical instrument for evaluating burn depth. *IEEE Trans Biomed Eng* 1988;35:842-50.
119. Green M, Holloway GA, Heimbach DM. Laser Doppler monitoring of microcirculatory changes in acute burn wounds. *J Burn Care Rehabil* 1988;9:57-62.
120. Cole RP, Shakespeare PG, Chissell HG, Jones SG. Thermographic assessment of burns using a nonpermeable membrane as wound covering. *Burns* 1991;17:117-22.
121. Wyllie FJ, Sutherland AB. Measurement of surface temperature as an aid to the diagnosis of burn depth. *Burns* 1991;17:123-7.
122. Cole RP, Jones SG, Shakespeare PG. Thermographic assessment of hand burns. *Burns* 1990;16:60-3.
123. Wachtel TL, Leopold GR, Frank HA, Frank DH. B-mode ultrasonic echo determination of depth of thermal injury. *Burns, including Thermal Injury* 1986;12:432-7.
124. Gatti JE, LaRossa D, Silverman DG, Hartford CE. Evaluation of the burn wound with perfusion fluorometry. *J Trauma* 1983;23:202-6.
125. Black KS, Hewitt CW, Miller DM, et al. Burn depth evaluation with fluorometry: Is it really definitive? *J Burn Care Rehabil* 1986;7:313-7.
126. Zawacki BE, Walker HL. An evaluation of patent blue V, bromphenol blue, and tetracycline for the diagnosis of burn depth. *Plast Reconstr Surg* 1970;45:459-65.
127. Chvapil M, Speer DP, Owen JA, Chvapil TA. Identification of the depth of burn injury by collagen stainability. *Plast Reconstr Surg* 1984;73:438-41.
128. Koruda MJ, Zimbler A, Settle RG, et al. Assessing burn wound depth using in vitro nuclear magnetic resonance (NMR). *J Surg Res* 1986;40:475-81.
129. Schweizer MP, Olsen JI, Shelby J, et al. Noninvasive assessment of metabolism in wounded skin by <sup>31</sup>P-NMR in vivo. *J Trauma* 1992;33:828-34.
130. Sheridan RL, Schomaker KT, Lucchina LC, et al. Burn depth estimation by use of indocyanine green fluorescence: initial human trial. *J Burn Care Rehabil* 1995;16:602-4.
131. Sheridan RL, Petras L, Basha G, et al. Planimetry study of the percent of body surface represented by the hand and palm: sizing irregular burns is more accurately done with the palm. *J Burn Care Rehabil* 1995;16:605-6.
132. Desai MH, Rutan RL, Hemdon DN. Conservative treatment of scald burns is superior to early excision. *J Burn Care Rehabil* 1991;12:482-4.
133. McDonald WS, Deitch EA. Hypertrophic skin grafts in burned patients: a prospective analysis of variables. *J Trauma* 1987;27:147-50.
134. Deitch EA, Wheelahan TM, Rose MP, Clothier J, Cotter J. Hypertrophic burn scars: analysis of variables. *J Trauma* 1983;23:895-8.

135. Soroff HS, Sasvary DH. Collagenase ointment and polymyxin B sulfate/bacitracin spray versus silver sulfadiazine cream in partial-thickness burns: a pilot study. *J Burn Care Rehabil* 1994;15:13-7.
136. Brown RA, Grobbelaar AO, Barker S, Rode H. A formula to calculate blood cross-match requirements for early burn surgery in children. *Burns* 1995;21:371-3.
137. Dye DJ. Requirements for cross-matched blood in burns surgery. *Burns* 1993;19:524-8.
138. Budny PG, Regan PJ, Roberts AH. The estimation of blood loss during burns surgery. *Burns* 1993;19:134-7.
139. Moran KT, O'Reilly TJ, Furman W, Munster AM. A new algorithm for calculation of blood loss in excisional burn surgery. *Am Surg* 1988;54:207-8.
140. Sheridan RL, Szyfelbein SK. Trends in blood conservation in burn care. *Burns* 2001;27:272-6.
141. Sheridan RL, Szyfelbein SK. Staged high dose epinephrine clays in pediatric burn excisions. *Burns* 1999;25:745-8.
142. Sheridan RL, Behringer GE, Ryan CM, et al. Effective postoperative protection for grafted posterior surfaces: the quilted dressing. *J Burn Care Rehabil* 1995;16:607-9.
143. Heimbach DM, Engrav LH, Marvin JA, Harnar TJ, Grube BJ. Toxic epidermal necrolysis: a step forward in treatment [published erratum appears in *JAMA* 1987;258:1894]. *JAMA* 1987;257:2171-5.
144. Fine JD, Johnson LB, Cronce D, et al. Intracytoplasmic retention of type VII collagen and dominant dystrophic epidermolysis bullosa: reversal of defect following cessation of or marked improvement in disease activity. *J Invest Dermatol* 1993;101:232-6.
145. Sheridan RL, Tompkins RG. Skin substitutes in burns. *Burns* 1999;25:97-103.
146. Song IC, Bromberg BE, Mohn MP, Koehnlein E. Heterografts as biological dressings for large skin wounds. *Surgery* 1966;59:576-83.
147. Elliott RA Jr, Hoehn JG. Use of commercial porcine skin for wound dressings. *Plast Reconstr Surg* 1973;52:401-5.
148. Ersek RA, Hachen HJ. Porcine xenografts in the treatment of pressure ulcers. *Ann Plast Surg* 1980;5:464-70.
149. Chatterjee DS. A controlled comparative study of the use of porcine xenograft in the treatment of partial thickness skin loss in an occupational health centre. *Curr Med Res Opin* 1978;5:726-33.
150. Marvin JA, Heimbach DM, Engrav LH, Harnar TJ. Improved treatment of the Stevens-Johnson syndrome. *Arch Surg* 1984;119:601-5.
151. Salisbury RE, Wilmore DW, Silverstein P, Pruitt BA Jr. Biological dressings for skin graft donor sites. *Arch Surg* 1973;106:705-6.
152. Salisbury RE, Carnes RW, Enterline D. Biological dressings and evaporative water loss from burn wounds. *Ann Plast Surg* 1980;5:270-2.
153. Demling RH. Burns. *N Engl J Med* 1985;313:1389-98.
154. Vanstraelen P. Comparison of calcium sodium alginate (KALTOSTAT) and porcine xenograft (E-Z DERM) in the healing of split-thickness skin graft donor sites. *Burns* 1992;18:145-8.
155. Bacha EA, Sheridan RL, Donohue GA, Tompkins RG. Staphylococcal toxic shock syndrome in a paediatric burn unit. *Burns* 1994;20:499-502.
156. Greenhalgh DG. The role of growth factors in wound healing. *J Trauma* 1996;41:159-67.

157. Bennett NT, Schultz GS. Growth factors and wound healing: biochemical properties of growth factors and their receptors. *Am J Surg* 1993;165:728-37.
158. Phillips TJ, Gilchrest BA. Cultured epidermal allografts as biological wound dressings. *Prog Clin Biol Res* 2000;365:77-94.
159. Teepe RG, Koch R, Haeseker B. Randomized trial comparing cryopreserved cultured epidermal allografts with tulle-gras in the treatment of split-thickness skin graft donor sites. *J Trauma* 1993;35:850-4.
160. Salzberg CA, Norris J, Carr J. Use of a biological dressing for the early treatment of deep partial thickness burns [abstract]. *J Burn Care Rehabil: Proceedings of the American Burn Association*. 1997;18:S102.
161. Miller MS. Use of topical recombinant human platelet-derived growth factor-BB (becaplermin) in healing of chronic mixed arteriovenous lower extremity diabetic ulcers. *J Foot Ankle Surg* 2000;38:227-31.
162. Hansbrough JF, Morgan J, Greenleaf G, Underwood J. Development of a temporary living skin replacement composed of human neonatal fibroblasts cultured in Biobrane, a synthetic dressing material. *Surgery* 1994;115:633-44.
163. Hansbrough J. Dermagraft-TC for partial-thickness burns: a clinical evaluation. *J Burn Care Rehabil* 1997;18:S25-8.
164. Parente ST. Estimating the economic cost offsets of using Dermagraft-TC as an alternative to cadaver allograft in the treatment of graftable burns. *J Burn Care Rehabil* 1997;18:S18-24.
165. Spielvogel RL. A histological study of Dermagraft-TC in patients' burn wounds. *J Burn Care Rehabil* 1997;18:S16-8.
166. Purdue GF. Dermagraft-TC pivotal efficacy and safety study. *J Burn Care Rehabil* 1997;18:S13-4.
167. Pape SA, Byrne PO. Safety and efficacy of transcyte for the treatment of partial-thickness burns. *J Burn Care Rehabil* 2000;21:390.
168. Noordenbos J, Dore C, Hansbrough JF. Safety and efficacy of TransCyte for the treatment of partial-thickness burns. *J Burn Care Rehabil* 2000;20:275-81.
169. Demling RH, DeSanti L. Management of partial thickness facial burns (comparison of topical antibiotics and bio-engineered skin substitutes). *Burns* 2000;25:256-61.
170. Morgan JR, Barrandon Y, Green H, Mulligan RC. Expression of an exogenous growth hormone gene by transplantable human epidermal cells. *Science* 1987;237:1476-9.
171. Sheridan RL, Morgan JR, Mohammed R. Polymeric biomaterials in burns and wounds. In: Sewerian D, editor. *Polymeric biomaterials*. New York: Marcel Dekker; In press.
172. Ramakrishnan KM, Jayaraman V. Management of partial-thickness burn wounds by amniotic membrane: a cost-effective treatment in developing countries. *Burns* 1997;23(suppl):S33-6.
173. Subrahmanyam M. Amniotic membrane as a cover for microskin grafts. *Br J Plast Surg* 1995;48:477-8.
174. Ganatra MA, Durrani KM. Method of obtaining and preparation of fresh human amniotic membrane for clinical use. *Journal of the Pakistan Medical Association* 1996;46:126-8.
175. Thomson PD, Parks DH. Monitoring, banking, and clinical use of amnion as a burn wound dressing. *Ann Plast Surg* 1981;7:354-6.
176. Quinby WC Jr, Hoover HC, Schefflan M, Walters PT, Slavin SA, Bondoc CC.

- Clinical trials of amniotic membranes in burn wound care. *Plast Reconstr Surg* 1982;70:711-7.
177. Bondoc CC, Burke JF. Clinical experience with viable frozen human skin and a frozen skin bank. *Ann Surg* 1971;174:371-82.
  178. Spence RJ. The banking and clinical use of human skin allograft in trauma patients: history. *Maryland Med J* 1986;35:50-2.
  179. May SR, Still JM Jr, Atkinson WB. Recent developments in skin banking and the clinical uses of cryopreserved skin. *J Med Assoc Ga* 1957;73:233-6.
  180. Burke JF, May JW Jr, Albright N, Quinby WC, Russell PS. Temporary skin transplantation and immunosuppression for extensive burns. *N Engl J Med* 1974;290:269-71.
  181. Bale JF Jr, Kealey GP, Ebelhack CL, Platz CE, Goeken JA. Cytomegalovirus infection in a cyclosporine-treated burn patient: case report. *J Trauma* 1992;32:263-7.
  182. Rheinwald JG, Green H. Serial cultivation of strains of human epidermal keratinocytes: the formation of keratinizing colonies from single cells. *Cell* 1975;6:331-43.
  183. Green H, Kehinde O, Thomas J. Growth of cultured human epidermal cells into multiple epithelia suitable for grafting. *Proc Natl Acad Sci U S A* 1979;76:5665-8.
  184. Green H. Cultured cells for the treatment of disease. *Sci Am* 1991;265:96-102.
  185. Gallico GG 3d, O'Connor NE, Compton CC, Kehinde O, Green H. Permanent coverage of large burn wounds with autologous cultured human epithelium. *N Engl J Med* 1984;311:448-51.
  186. Gallico GG 3d, O'Connor NE, Compton CC, Remensnyder JP, Kehinde O, Green H. Cultured epithelial autografts for giant congenital nevi. *Plast Reconstr Surg* 1989;84:1-9.
  187. Sheridan RL, Tompkins RG. Cultured autologous epithelium in patients with burns of ninety percent or more of the body surface. *J Trauma* 1995;38:48-50.
  188. Rue LW III, Cioffi WG, McManus WF, Pruitt BA Jr. Wound closure and outcome in extensively burned patients treated with cultured autologous keratinocytes. *J Trauma* 1993;34:662-7.
  189. Langdon RC, Cuono CB, Birchall N, et al. Reconstitution of structure and cell function in human skin grafts derived from cryopreserved allogeneic dermis and autologous cultured keratinocytes. *J Invest Dermatol* 1988;91:478-85.
  190. Cuono CB, Langdon R, Birchall N, Bartelbort S, McGuire J. Composite autologous-allogeneic skin replacement: development and clinical application. *Plast Reconstr Surg* 1987;80:626-37.
  191. Cuono C, Langdon R, McGuire J. Use of cultured epidermal autografts and dermal allografts as skin replacement after burn injury. *Lancet* 1986;1:1123-4.
  192. Hickerson WL, Compton C, Fletchall S, Smith LR. Cultured epidermal autografts and allodermis combination for permanent burn wound coverage. *Burns* 1994;20(suppl):S52-6.
  193. Compton CC, Hickerson W, Nadire K, Press W. Acceleration of skin regeneration from cultured epithelial autografts by transplantation to homograft dermis. *J Burn Care Rehabil* 1993;14:653-62.
  194. Tompkins RG, Burke JF. Progress in burn treatment and the use of artificial skin. *World J Surg* 1990;14:819-24.
  195. Yannas IV, Burke JF, Warpehoski M, et al. Prompt, long-term functional replacement of skin. *Transact Am Soc Artif Internal Organs* 1981;27:19-23.

196. Yannas IV, Burke JF, Orgill DP, Skrabut EM. Wound tissue can utilize a polymeric template to synthesize a functional extension of skin. *Science* 1982;215:174-6.
197. Tompkins RG, Hilton JF, Burke JF, et al. Increased survival after massive thermal injuries in adults: preliminary report using artificial skin. *Crit Care Med* 1989;17:734-40.
198. Scalea TM, Simon HM, Duncan AO, et al. Geriatric blunt multiple trauma: improved survival with early invasive monitoring. *J Trauma* 1990;30:129-36.
199. Sheridan RL, Heggerty M, Tompkins RG, Burke JF. Artificial skin in massive burns: results at ten years. *Eur J Plast Surg* 1994;17:91-3.
200. Heimbach D, Luterman A, Burke J, et al. Artificial dermis for major burns: a multi-center randomized clinical trial. *Ann Surg* 1988;208:313-20.
201. Wainwright DJ. Use of an acellular allograft dermal matrix (Alloderm) in the management of full-thickness burns. *Burns* 1995;21:243-8.
202. Wainwright D, Madden M, Luterman A, et al. Clinical evaluation of an acellular allograft dermal matrix in full-thickness burns. *J Burn Care Rehabil* 1996;17:124-36.
203. Sheridan RL, Choucair RJ. Acellular allograft dermis does not hinder initial engraftment in burn resurfacing and reconstruction. *J Burn Care Rehabil* 1997;18:496-9.
204. Sheridan RL, Choucair RJ. Acellular allodermis in burn surgery: 1-year results of a pilot trial [abstract]. *J Burn Care Rehabil* 1998;19:528-30.
205. Cooper ML, Hansbrough JF, Spielvogel RL, Cohen R, Bartel RL, Naughton G. In vivo optimization of a living dermal substitute employing cultured human fibroblasts on a biodegradable polyglycolic acid or polyglactin mesh. *Biomaterials* 1991;12:243-8.
206. Hansbrough JF, Cooper ML, Cohen R, et al. Evaluation of a biodegradable matrix containing cultured human fibroblasts as a dermal replacement beneath meshed skin grafts on athymic mice. *Surgery* 1992;111:438-46.
207. Hansbrough JF, Dore C, Hansbrough WB. Clinical trials of a living dermal tissue replacement placed beneath meshed, split-thickness skin grafts on excised burn wounds. *J Burn Care Rehabil* 1992;13:519-29.
208. Briggaman RA, Wheeler CE Jr. The epidermal-dermal junction. *J Invest Dermatol* 1975;65:71-84.
209. Cooper ML, Andree C, Hansbrough JF, Zapata-Sirvent RL, Spielvogel RL. Direct comparison of a cultured composite skin substitute containing human keratinocytes and fibroblasts to an epidermal sheet graft containing human keratinocytes on athymic mice. *J Invest Dermatol* 1993;101:811-9.
210. Woodley DT, Peterson HD, Herzog SR, et al. Burn wounds resurfaced by cultured epidermal autografts show abnormal reconstitution of anchoring fibrils. *JAMA* 1988;259:2566-71.
211. Coulomb B, Lebreton C, Dubertret L. Influence of human dermal fibroblasts on epidermalization. *J Invest Dermatol* 1989;92:122-5.
212. Boyce ST, Christianson DJ, Hansbrough JF. Structure of a collagen-GAG dermal skin substitute optimized for cultured human epidermal keratinocytes. *J Biomed Mater Res* 1988;22:939-57.
213. Boyce ST, Hansbrough JF. Biologic attachment, growth, and differentiation of cultured human epidermal keratinocytes on a graftable collagen and chondroitin-6-sulfate substrate. *Surgery* 1988;103:421-31.
214. Cooper ML, Hansbrough JF. Use of a composite skin graft composed of cultured

- human keratinocytes and fibroblasts and a collagen-GAG matrix to cover full-thickness wounds on athymic mice. *Surgery* 1991;109:198-207.
215. Hansbrough JF, Boyce ST, Cooper ML, Foreman TJ. Burn wound closure with cultured autologous keratinocytes and fibroblasts attached to a collagen-glycosaminoglycan substrate. *JAMA* 1989;262:2125-30.
  216. Hansbrough JF, Morgan JL, Greenleaf GE, Bartel R. Composite grafts of human keratinocytes grown on a polyglactin mesh-cultured fibroblast dermal substitute function as a bilayer skin replacement in full-thickness wounds on athymic mice. *J Burn Care Rehabil* 1993;14:485-94.
  217. Sheridan R, Morgan JR, Cusick J, Petras L, Lydon M, Tompkins R. Initial experience with an autologous composite skin replacement. *Burns* 2001;27:421-4.
  218. Compton CC, Butler CE, Yannas IV, Warland G, Orgill DP. Organized skin structure is regenerated in vivo from collagen-GAG matrices seeded with autologous keratinocytes. *J Invest Dermatol* 2000;110:908-16.
  219. Brues AM. The pulmonary complications: a clinical description. *Ann Surg* 1943;117:834-40.
  220. Hunt JL, Agee RN, Pruitt BA Jr. Fiberoptic bronchoscopy in acute inhalation injury. *J Trauma* 1975;15:641-9.
  221. Lin WY, Kao CH, Wang SJ. Detection of acute inhalation injury in fire victims by means of technetium-99m DTPA radioaerosol inhalation lung scintigraphy. *Eur J Nucl Med* 1997;24:125-9.
  222. Lull RJ, Anderson JH, Telepak RJ, Brown JM, Utz JA. Radionuclide imaging in the assessment of lung injury. *Semin Nucl Med* 1980;10:302-10.
  223. Parker JC, Hernandez LA, Peevy KJ. Mechanisms of ventilator-induced lung injury. *Crit Care Med* 1993;21:131-43.
  224. Desai MH, Mlcak RP, Robinson E, et al. Does inhalation injury limit exercise endurance in children convalescing from thermal injury? *J Burn Care Rehabil* 1993;14:12-6.
  225. Weiss LD, Van Meter KW. The applications of hyperbaric oxygen therapy in emergency medicine. *Am J Emerg Med* 1992;10:558-68.
  226. Thom SR, Taber RL, Mendiguren II, Clark JM, Hardy KR, Fisher AB. Delayed neuropsychologic sequelae after carbon monoxide poisoning: prevention by treatment with hyperbaric oxygen. *Ann Emerg Med* 1995;25:474-80.
  227. Myers RA, Snyder SK, Emhoff TA. Subacute sequelae of carbon monoxide poisoning. *Ann Emerg Med* 1985;14:1163-7.
  228. Sheridan RL. Hyperbaric oxygen treatment: a brief overview of a controversial topic. *J Trauma* 1999;47:426-35.
  229. Scheinkestel CD, Bailey M, Myles PS, et al. Hyperbaric or normobaric oxygen for acute carbon monoxide poisoning: a randomised controlled clinical trial. *Med J Aust* 2000;170:203-10.
  230. Grube BJ, Marvin JA, Heimbach DM. Therapeutic hyperbaric oxygen: help or hindrance in burn patients with carbon monoxide poisoning? *J Burn Care Rehabil* 1988;9:249-52.
  231. Barillo DJ, Goode R, Esch V. Cyanide poisoning in victims of fire: analysis of 364 cases and review of the literature. *J Burn Care Rehabil* 1994;15:46-57.
  232. Sheridan RL. Airway management and respiratory care of the burn patient. *Int Anesthesiol Clin* 2000;38:129-45.

233. Fitzpatrick JC, Cioffi WG Jr, Cheu HW, Pruitt BA Jr. Predicting ventilation failure in children with inhalation injury. *J Pediatr Surg* 1994;29:1122-6.
234. Niederman MS, Torres A, Summer W. Invasive diagnostic testing is not needed routinely to manage suspected ventilator-associated pneumonia. *Am J Respir Crit Care Med* 1994;150:565-9.
235. Hollingsed TC, Saffle JR, Barton RG, Craft WB, Morris SE. Etiology and consequences of respiratory failure in thermally injured patients. *Am J Surg* 1993;166:592-6.
236. Slutsky AS. Mechanical ventilation: American College of Chest Physicians' Consensus Conference. *Chest* 1993;104:1833-59.
237. Shapiro BA, Cane RD, Harrison RA, Steiner MC. Changes in intrapulmonary shunting with administration of 100 percent oxygen. *Chest* 1980;77:138-41.
238. Cane RD, Peruzzi WT, Shapiro BA. Airway pressure release ventilation in severe acute respiratory failure. *Chest* 1991;100:460-3.
239. Webb HH, Tierney DF. Experimental pulmonary edema due to intermittent positive pressure ventilation with high inflation pressures: protection by positive end-expiratory pressure. *Am Rev Respir Dis* 1974;110:556-65.
240. Sheridan RL, Kacmarek RM, McEttrick MM, et al. Permissive hypercapnia as a ventilatory strategy in burned children: effect on barotrauma, pneumonia, and mortality. *J Trauma* 1995;39:854-9.
241. Hickling KG, Walsh J, Henderson S, Jackson R. Low mortality rate in adult respiratory distress syndrome using low-volume, pressure-limited ventilation with permissive hypercapnia: a prospective study. *Crit Care Med* 1994;22:1568-78.
242. Hickling KG, Downward G, Davis FM, A'Court G. Management of severe ARDS with low frequency positive pressure ventilation and extracorporeal CO<sub>2</sub> removal. *Anaesth Intensive Care* 1986;14:79-83.
243. Sheridan RL, Hurford WE, Kacmarek RM, et al. Inhaled nitric oxide in burn patients with respiratory failure. *J Trauma* 1997;42:629-34.
244. Sheridan RL, Zapol WM, Ritz RH, et al. Low-dose inhaled nitric oxide in acutely burned children with severe respiratory failure. *Surgery* 1999;126:856-62.
245. Rodeberg DA, Housinger TA, Greenhalgh DG, Maschinot NE, Warden GD. Improved ventilatory function in burn patients using volumetric diffusive respiration. *J Am Coll Surg* 1994;179:518-22.
246. Goretsky MJ, Greenhalgh DG, Warden GD, Ryckman FC, Warner BW. The use of extracorporeal life support in pediatric burn patients with respiratory failure. *J Pediatr Surg* 1995;30:620-3.
247. Sellers BJ, Davis BL, Larkin PW, Morris SE, Saffle JR. Early prediction of prolonged ventilator dependence in thermally injured patients. *J Trauma* 1997;43:899-903.
248. Jones WG, Madden M, Finkelstein J, Yurt RW, Goodwin CW. Tracheostomies in burn patients. *Ann Surg* 1989;209:471-4.
249. Hunt JL, Purdue GF, Gunning T. Is tracheostomy warranted in the burn patient? Indications and complications. *J Burn Care Rehabil* 1986;7:492-5.
250. Calhoun KH, Deskin RW, Garza C, et al. Long-term airway sequelae in a pediatric burn population. *Laryngoscope* 1988;98:721-5.
251. Anene O, Meert KL, Uy H, Simpson P, Sarnaik AP. Dexamethasone for the prevention of postextubation airway obstruction: a prospective, randomized, double-blind, placebo-controlled trial. *Crit Care Med* 1996;24:1666-9.

252. NIH Consensus Statement: NIH Consensus Statement. *Crit Care Med* 1983;46:1-26.
253. Sheridan RL. Critical care of the burn patient. In: Hurford WE, Bigatello LM, Haspel KL, Warren RL, editors. *Critical Care Handbook of the Massachusetts General Hospital*. Philadelphia: Lippincott, Williams, and Wilkins; 2000. p. 584-609.
254. Marquez S, Turley JJ, Peters WJ. Neuropathy in burn patients. *Brain* 1993;116:471-83.
255. Dagum AB, Peters WJ, Neligan PC, Douglas LG. Severe multiple mononeuropathy in patients with major thermal burns. *J Burn Care Rehabil* 1993;14:440-5.
256. Stoddard FM, Martyn JJ, Sheridan RL. Psychiatric issues in pain of burn injury: controlling pain and improving outcomes. *Curr Rev Pain*. 1997;1:130-6.
257. Sheridan RL, McEtrick M, Bacha G, Stoddard F, Tompkins RG. Midazolam infusion in pediatric patients with burns who are undergoing mechanical ventilation. *J Burn Care Rehabil* 1994;15:515-8.
258. Sheridan R, Stoddard F, Querzoli E. Management of background pain and anxiety in critically burned children requiring protracted mechanical ventilation. *J Burn Care Rehabil* 2001;22:150-3.
259. Bondoc CC, Morris PJ, Wee T, Burke JF. Metabolic effects of 0.5 per cent silver nitrate therapy for extensive burns in children. *Surg Forum* 1966;17:475-7.
260. Burke JF, Bondoc CC, Morris PJ. Metabolic effects of topical silver nitrate therapy in burns covering more than fifteen percent of the body surface. *Ann N Y Acad Sci* 1968;150:674-80.
261. Zaloga GP. Interpretation of the serum magnesium level [editorial]. *Chest* 1989;95:257-8.
262. Wu JZ, Ogle CK, Fischer JE, Warden GD, Ogle JD. The mRNA expression and in vitro production of cytokines and other proteins by hepatocytes and Kupffer cells following thermal injury. *Shock* 1995;3:268-73.
263. Greenhalgh DG, Housinger TA, Kagan RJ, et al. Maintenance of serum albumin levels in pediatric burn patients: a prospective, randomized trial. *J Trauma* 1995;39:67-74.
264. Guenter PA, Settle RG, Perlmutter S, Marino PL, DeSimone GA, Rolandelli RH. Tube feeding-related diarrhea in acutely ill patients. *JPEN: J Parenter Enter Nutr* 1991;15:277-80.
265. Sapijaszko MJ, Brant R, Sandham D, Berthiaume Y. Nonrespiratory predictor of mechanical ventilation dependency in intensive care unit patients. *Crit Care Med* 1996;24:601-7.
266. Sheridan RL, Prelack K, Cunningham JJ. Physiologic hypoalbuminemia is well tolerated by severely burned children. *J Trauma* 1997;43:448-52.
267. Sheridan RL, Weber JM, Peterson HF, Tompkins RG. Central venous catheter sepsis with weekly catheter change in paediatric burn patients: an analysis of 221 catheters. *Burns* 1995;21:127-9.
268. Goldstein AM, Weber JM, Sheridan RL. Femoral venous access is safe in burned children: an analysis of 224 catheters. *J Pediatr* 1997;130:442-6.
269. Strome DR, Aulick LH, Mason AD Jr, Pruitt BA Jr. Thermoregulatory and non-thermoregulatory heat production in burned rat. *J Appl Physiol* 1986;61:688-93.
270. Souba WW. Cytokine control of nutrition and metabolism in critical illness. *Curr Probl Surg* 1994;31:577-643.
271. Carlson DE, Cioffi WG Jr, Mason AD Jr, McManus WF, Pruitt BA Jr. Resting

- energy expenditure in patients with thermal injuries. *Surg Gynecol Obstet* 1992; 174:270-6.
272. Sheridan RL, Prelack K, Yin L. Energy needs are poorly predicted in the critically ill elderly. *J Int Care Med* 1997;12:45-9.
  273. Saffle JR, Medina E, Raymond J, Westenskow D, Kravitz M, Warden GD. Use of indirect calorimetry in the nutritional management of burned patients. *J Trauma* 1985;25:32-9.
  274. Prelack K, Cunningham JJ, Sheridan RL, Tompkins RG. Energy and protein provisions for thermally injured children revisited: an outcome-based approach for determining requirements. *J Burn Care Rehabil* 1997;18:177-81.
  275. Gilpin DA, Barrow RE, Rutan RL, Broemeling L, Herndon DN. Recombinant human growth hormone accelerates wound healing in children with large cutaneous burns. *Ann Surg* 1994;220:19-24.
  276. Meyer NA, Muller MJ, Herndon DN. Nutrient support of the healing wound. *New Horizons* 1994;2:202-14.
  277. Martineau L, Little RA, Rothwell NJ, et al. Clenbuterol, a beta 2-adrenergic agonist, reverses muscle wasting due to scald injury in the rat: the acute-phase response of cultured rat hepatocytes: system characterization and the effect of human cytokines. *Burns* 1984;224:505-14.
  278. Mendenhall CL, Moritz TE, Roselle GA, et al. A study of oral nutritional support with oxandrolone in malnourished patients with alcoholic hepatitis: results of a Department of Veterans Affairs cooperative study Investigations on the influence of Clenbuterol and recombinant porcine somatotropin on nitrogen and energy metabolism in growing pigs. *Hepatology* 1993;45:1-11.
  279. Mainous MR, Block EF, Deitch EA. Nutritional support of the gut: how and why. *New Horizons*. 1994;2:193-201.
  280. Hu OY, Ho ST, Wang JJ, Ho W, Wang HJ, Lin CY. Evaluation of gastric emptying in severe, burn-injured patients. *Crit Care Med* 1993;21:527-31.
  281. Kadillack P, Prelack K, Sheridan R. Gastric tube feedings with supplemental parenteral nutrition in children with large burns. *J Burn Care Rehabil* 1999;20:S248.
  282. Weber JM, Sheridan RL, Pasternack MS, Tompkins RG. Nosocomial infections in pediatric patients with burns. *Am J Infect Control* 1997;25:195-201.
  283. Sheridan RL, Weber JM, Tompkins RG. Candidemia in pediatric burn patient. *J Burn Care Rehabil* 1995;16:440-3.
  284. Sheridan RL, Ryan CM, Tompkins RG. Acute adrenal insufficiency in the burn intensive care unit. *Burns* 1993;19:63-6.
  285. Sheridan RL, Ryan CM, Pasternack MS, Weber JM, Tompkins RG. Flavobacterial sepsis in massively burned pediatric patients. *Clin Infect Dis* 1993;17:185-7.
  286. Sheridan RL, Weber J, Benjamin J, Pasternack MS, Tompkins RG. Control of methicillin-resistant *Staphylococcus aureus* in a pediatric burn unit. *Am J Infect Control* 1994;22:340-5.
  287. Goodwin CW. Multiple organ failure: clinical overview of the syndrome. *J Trauma* 1990;30:S163-5.
  288. Gladston I. *Progress in medicine*. New York: Knopf; 1940. p. 292.
  289. Sheridan RL, Szyfelbein SK. Staged high-dose epinephrine claysis is safe and effective in extensive tangential burn excisions in children. *Burns* 2000;25:745-8.
  290. Asch MJ, Moylan JA Jr, Bruck HM, Pruitt BA Jr. Ocular complications associated with burns: review of a five-year experience including 104 patients. *J Trauma* 1971; 11:857-61.

291. Sheridan RL, Hurley J, Smith MA, et al. The acutely burned hand: management and outcome based on a ten-year experience with 1047 acute hand burns. *J Trauma* 1995;38:406-11.
292. Sheridan RL, Baryza MJ, Pessina MA, et al. Acute hand burns in children: management and long-term outcome based on a 10-year experience with 698 injured hands. *Ann Surg* 1999;229:558-64.
293. Housinger TA, Keller B, Warden GD. Management of burns of the penis. *J Burn Care Rehabil* 1993;14:525-7.
294. Sheridan RL, Gagnon SW, Tompkins RG. The burn unit as a resource for the management of acute nonburn conditions in children. *J Burn Care Rehabil* 1995;16:62-4.
295. Rabban JT, Blair JA, Rosen CL, Adler JN, Sheridan RL. Mechanisms of pediatric electrical injury: new implications for product safety and injury prevention. *Arch Pediatr Adolesc Med* 1997;151:696-700.
296. Donelan MB. Reconstruction of electrical burns of the oral commissure with a ventral tongue flap. *Plast Reconstr Surg* 1995;95:1155-64.
297. Mann R, Gibran N, Engrav L, Heimbach D. Is immediate decompression of high voltage electrical injuries to the upper extremity always necessary? *J Trauma* 1996;40:584-9.
298. Sheridan RL, Ryan CM, Quinby WC Jr, Blair J, Tompkins RG, Burke JF. Emergency management of major hydrofluoric acid exposures. *Burns* 1995;21:62-4.
299. Stratta RJ, Saffle JR, Kravitz M, Warden GD. Management of tar and asphalt injuries. *Am J Surg* 1983;146:766-9.
300. Demling RH, Buerstatte WR, Perea A. Management of hot tar burns. *J Trauma* 1980;20:242.
301. Britt LD, Dascombe WH, Rodriguez A. New horizons in the management of hypothermia and frostbite injury. *Surg Clin North Am* 1991;71:345-70.
302. Pruitt BA Jr. Burn treatment for the unburned [editorial]. *JAMA* 1987;257:2207-8.
303. Sheridan RL, Weber JM, Schulz JT, Ryan CM, Low HM, Tompkins RG. Management of severe toxic epidermal necrolysis in children. *J Burn Care Rehabil* 2000;20:497-500.
304. Sheridan RL, Schulz JT, Ryan CM, et al. Long-term consequences of toxic epidermal necrolysis in children. *Pediatrics* (In Press).
305. Kelemen JJ 3rd, Cioffi WG, McManus WF, Mason AD Jr, Pruitt BA Jr. Burn center care for patients with toxic epidermal necrolysis. *J Am Coll Surg* 1995;180:273-8.
306. Adzick NS, Kim SH, Bondoc CC, Quinby WC, Remensnyder JP. Management of toxic epidermal necrolysis in a pediatric burn center. *Am J Dis Child* 1985;139:499-502.
307. Schulz JT, Sheridan RL, Ryan CM, MacKool B, Tompkins RG. A 10-year experience with toxic epidermal necrolysis. *J Burn Care Rehabil* 2000;21:199-204.
308. Sheridan RL, Briggs SE, Remensnyder JP, Tompkins RG. Management strategy in purpura fulminans with multiple organ failure in children. *Burns* 1996;22:53-6.
309. Aiuto LT, Barone SR, Cohen PS, Boxer RA. Recombinant tissue plasminogen activator restores perfusion in meningococcal purpura fulminans. *Crit Care Med* 1997;25:1079-82.
310. Sheridan RL, Briggs SE, Remensnyder JP, et al. The burn unit as a resource for the management of acute nonburn conditions in children. *J Burn Care Rehabil* 1995;16:62-4.
311. Sheridan RL, Weber JM, Pasternack MM, Tompkins RG. Admission streptococcal screening allows elimination of burn wound prophylaxis. *J Trauma*. In press.
312. Payne JH. Home Sweet Home from Clari, the Maid of Milan. In: Turner MR, editor.

- Favorite parlour poetry—an annotated anthology. New York: Dover Publications; 1992. p. 325.
313. Coffee T, Yurko L, Fratianne RB. Mixing inpatient with outpatient care: establishing an outpatient clinic on a burn unit. *J Burn Care Rehabil* 1992;13:587-9.
  314. Waitzman AA, Neligan PC. How to manage burns in primary care. *Can Family Phys* 1993;39:2394-400.
  315. Heinrich JJ, Brand DA, Cuono CB. The role of topical treatment as a determinant of infection in outpatient burns. *J Burn Care Rehabil* 1988;9:253-7.
  316. Salisbury R. Burn rehabilitation: our unanswered challenge: the 1992 presidential address to the American Burn Association. *J Burn Care Rehabil* 1992;13:495-505.
  317. Kischer CW. The microvessels in hypertrophic scars, keloids and related lesions. *J Submicrosc Cytol Path* 1992;24:281-96.
  318. Beranek JT, Masseyeff R. Hyperplastic capillaries and their possible involvement in the pathogenesis of fibrosis. *Histopathology* 1992;10:543-51.
  319. Sheridan RL, MacMillan K, Donelan M, et al. Tunable dye laser neovessel ablation as an adjunct to the management of hypertrophic scarring in burned children: pilot trial to establish safety. *J Burn Care Rehabil* 1997;18:317-20.
  320. Carr-Collins JA. Pressure techniques for the prevention of hypertrophic scar. *Clin Plast Surg* 1992;19:733-43.
  321. Clark JA, Cheng JC, Leung KS, Leung PC. Mechanical characterisation of human postburn hypertrophic skin during pressure therapy. *J Biomech* 1987;20:397-406.
  322. Ahn ST, Monafó WW, Mustoe TA. Topical silicone gel for the prevention and treatment of hypertrophic scar. *Arch Surg* 1991;126:499-504.
  323. Ridgway CL, Daugherty MB, Warden GD. Serial casting as a technique to correct burn scar contractures: a case report. *J Burn Care Rehabil* 1991;12:67-72.
  324. Achauer BM. Burn reconstruction. New York: Thieme Medical; 1991. p. 1-188.
  325. Peterson SL, Mani MM, Crawford CM, Neff JR, Hiebert JM. Postburn heterotopic ossification: insights for management decision making. *J Trauma* 1989;29:365-9.
  326. Richards AM, Klaassen MF. Heterotopic ossification after severe burns: a report of three cases and review of the literature. *Burns* 1997;23:64-8.
  327. Djurickovic S, Meek RN, Snelling CF, et al. Range of motion and complications after postburn heterotopic bone excision about the elbow. *J Trauma* 1996;41:825-30.
  328. O'Nan S. The circus fire. New York: Doubleday; 2000. p. 363.
  329. Stoddard FJ, Stroud L, Murphy JM. Depression in children after recovery from severe burns. *J Burn Care Rehabil* 1992;13:340-7.
  330. Rizzone LP, Stoddard FJ, Murphy JM, Kruger LJ. Posttraumatic stress disorder in mothers of children and adolescents with burns. *J Burn Care Rehabil* 1994;15:158-63.